

2021-22 North Carolina Avid Grouse Hunter Survey

Hannah Plumpton, *NCWRC Upland Game Bird Biologist* Mobile (984) 800-7222 hananh.plumpton@ncwildlife.org



Since 1984, the North Carolina Wildlife Resources Commission (NCWRC) has conducted an annual grouse hunter survey to estimate long-term grouse hunting trends and provide annual insight into grouse hunting demographics throughout the mountains of North Carolina. Volunteer grouse hunters participate by recording and submitting their annual hunting activity throughout the season. Grouse hunting activity is recorded by county and landownership type (Private Land or Game Land) within the two grouse management regions (Northern Mountains and Southern Mountains) (Fig. 1). Reported hunting trips typically consist of a single day per hunting party.




Figure 1. Grouse Management Regions and Ranger Districts on Pisgah and Nantahala National Forests in Western North Carolina.

Fifty-two avid grouse hunters reported information during the 2021-22 season, providing grouse hunting statistics for 416 hunting trips (Fig. 2). The gradual annual decline of total reported grouse hunting trips has primarily been a function of fewer hunters and fewer hunting trips per hunter. Presumably this is due to fewer grouse and poor hunting in recent years. Since the inception of the survey in 1984, Ashe and Madison Counties have had the most grouse hunts reported, with over 4,000 hunts occurring in each of these counties (Fig. 3). During the 2021-22 season, Macon and Clay Counties were most often reported, with more than 50 hunts reported from each.




Figure 2. Total number of reported hunts by volunteer avid grouse hunter survey participants, 1984-85 through 2021-22.




Figure 3. Total number of hunts by county as reported by volunteer avid grouse hunter survey participants, 1984-85 through 2021-22.

During the 2021-22 season, avid grouse survey participants hunted an average of 8 times (Fig. 4). It is clear that participants are now hunting considerably fewer times than during the 1980's and 1990's. The average length of a hunting trip has declined somewhat over that time period as well, with an average trip length of just 3.4 hours reported during the 2021-22 season (Fig. 5). This may be a result of aging hunters, fewer grouse, or a combination of both.




Figure 4. Average number of hunting trips per hunter based on avid grouse hunter survey participants, 1985-86 through 2021-22.




Figure 5. Average length (hours) of hunting trips of avid grouse hunter survey participants, 1984-85 through 2021-22.

Flush rates are presented both by hunting trip and by hours hunted in this report. Flush rates by hour may provide a more precise index to grouse abundance, while flush rates by hunting trip are more applicable from grouse hunting perspectives. However, we recognize that hunters will change their hunting locations over time to areas with relatively more grouse. This selective hunting behavior has a tendency to skew trend estimates such that they may not represent actual annual abundances or changes in abundance across the full landscape.

The avid grouse hunter survey has documented overall long-term declines in hourly flush rates. While some years have shown slight increases, the overall trend has been a steady decline. This has been true on both private land and Game Lands and in both the northern and southern mountain regions (Fig. 6). Historically more grouse were reported in the southern mountain region, however flush rates reported from the northern mountains have been very comparable for much of the last decade, and flush rates in both regions have declined to all-time lows (Fig. 7).




Figure 6. Average grouse flushed per hour by land type by avid grouse hunter survey participants, 1989-90 through 2021-22.




Figure 7. Average grouse flushed per hour by region by avid grouse hunter survey participants, 1984-85 through 2021-22.

Grouse hunting during the 2021-22 season was very poor in comparison to what hunters encountered when this survey began in the 1980's. In fact, grouse hunting this season was the one of the worst on record for the numbers of grouse flushed and bagged per trip (Figs. 8-9). On half (50%) of trips hunters did not flush any grouse (Fig. 10).




Figure 8. Average number of grouse flushed per hunting trip by avid grouse hunters, 1984-85 through 2021-22.




Figure 9. Average number of grouse bagged per hunting trip by avid grouse hunters, 1984-85 through 2021-22.




Figure 10. Percent of reported grouse hunting trips with no flushes by avid grouse hunters, 1984-85 through 2021-22.

Not surprisingly, during the 2021-22 hunting season, avid hunters reported more hunting activity later in the winter after big game hunting seasons have closed (Fig. 11). Grouse hunters made substantially more trips in January and February than earlier in the season. However, the number of grouse killed did not increase in a similar fashion. Hunters reported killing between 6 and 16 grouse each month of the season.




Figure 11. Total reported grouse hunting trips and harvests during the 2020-21 hunting season by avid grouse hunter survey participants.

West Nile Virus

During the past three hunting seasons (2018-19 through 2020-21) avid grouse hunters provided feathers and blood samples from 117 ruffed grouse. Laboratory tests revealed that 11% of the grouse were positive for West Nile virus antibodies. While it is unclear what impact West Nile virus may be having on the grouse population, it is very clear that exposure to this virus does occur at some level. The results tell us that some grouse killed by hunters were infected earlier in their life. It does not tell us when the infection occurred or whether the birds had any symptoms or health-related issues as a result of the virus. Also, it is important to realize that this does not tell us how many grouse may have died from infection and therefore were not available for hunters to harvest. Prevalence of 11% is fairly low, which could be a result of low levels of West Nile virus in the region, or it could mean that a high proportion of infected birds did not survive. We will continue our work here in North Carolina and will also continue to collaborate with our colleagues in other states to fully answer these questions.

Funding for the avid grouse hunter survey report was partially provided through a Pittman-Robertson Wildlife Restoration Multi-state Grant. The Federal Aid in Wildlife Restoration Act, popularly known as the Pittman-Robertson Act, was approved by Congress on September 2, 1937, and began functioning July 1, 1938. The purpose of this Act was to provide funding for the selection, restoration, rehabilitation and improvement of wildlife habitat, wildlife management research, and the distribution of information produced by the projects. The Act was amended October 23, 1970, to include funding for hunter training programs and the development, operation and maintenance of public target ranges.

Funds are derived from an 11 percent Federal excise tax on sporting arms, ammunition, and archery equipment, and a 10 percent tax on handguns. These funds are collected from the manufacturers by the Department of the Treasury and are apportioned each year to the States and Territorial areas (except Puerto Rico) by the Department of the Interior on the basis of formulas set forth in the Act. Funds for hunter education and target ranges are derived from one-half of the tax on handguns and archery equipment.

Each state's apportionment is determined by a formula which considers the total area of the state and the number of licensed hunters in the state. The program is a cost-reimbursement program, where the state covers the full amount of an approved project then applies for reimbursement through Federal Aid for up to 75 percent of the project expenses. The state must provide at least 25 percent of the project costs from a non-federal source





	% Hinto	No Grouse	Flushed	12.79%	14.17%	12.13%	11.39%	11.65%	8.82%	8.48%	10.53%	14.06%	15.20%	12.32%	15.41%	11.71%	12.54%	20.08%	19.13%	16.52%	14.73%	15.90%	17.95%	16.60%	18.48%	18.61%	25.39%	29.75%	29.23%	29.48%	20.00%	28.37%	28.57%	39.15%	25.99%	35.40%	47.81%	33.13%	43.00%	51.00%	49.51%
	# Hunts	Grouse	Flushed	172	140	104	109	126	111	109	131	197	176	192	198	193	182	239	246	204	174	152	152	132	122	131	177	213	161	194	203	261	238	330	183	245	306	168	193	230	506
		Grouse	Harvested/Hour	0.17	0.15	0.15	0.15	0.17	0.21	0.20	0.19	0.14	0.17	0.19	0.17	0.18	0.17	0.11	0.14	0.14	0.15	0.15	0.13	0.13	0.12	0.13	0.10	0.08	0.09	0.07	0.14	0.08	0.08	90'0	0.09	90.0	90'0	90'0	0.04	0.04	0.04
		Grouse	Harvested/Trip	69'0	99.0	09'0	0.64	0.71	0.91	0.89	0.81	0.58	0.66	0.78	29'0	0.70	69.0	0.43	0.55	0.55	0.59	0.59	0.51	05.0	0.47	0.52	0.36	0.30	0.33	0.28	0.51	0.33	0.30	0.23	0.33	0.22	0.20	0.22	0.13	0.12	0.14
	# 02102	Harvests	Reported	933	641	512	613	692	1151	1142	1008	817	692	1213	998	1152	1008	512	713	9/9	692	595	434	400	313	365	249	214	180	184	200	303	249	190	229	154	127	112	09	99	69
)-21 hi	o di cu	Ë							1.48	1.51	1.45	1.04	1.23	1.56	1.44	1.53	1.41	1.11	1.20	1.26	1.35	1.27	1.05	1.26	1.16	1.19	96.0	0.94	1.04	0.77	1.22	0.88	96'0	0.91	0.93	0.71	62'0	98'0	0.51	0.52	09.0
	o di locali	Flushed/Hour -	Game Land						1.20	1.27	1.05	0.94	0.90	1.08	0.74	1.15	0.94	0.62	0.89	0.79	0.82	0.88	0.81	0.74	0.79	0.88	0.52	0.48	0.50	09:0	0.94	0.54	0.58	0.35	0.63	0.50	0.28	69.0	0.41	0.35	0.43
	Grouse Flushad/Hour	Southern	Mountains	1.18	1.20	1.39	1.24	1.19	1.45	1.49	1.35	1.08	1.16	1.42	1.26	1.37	1.27	0.78	1.02	1.05	1.13	1.17	0.98	1.07	1.12	1.07	0.73	0.68	0.72	0.68	1.07	0.68	0.72	0.53	0.71	0.58	0.46	29.0	0.45	0.42	0.50
	Grouse Flushad/Hour	Northern	Mountains	92.0	92.0	0.84	06.0	1.15	1.19	1.27	1.17	0.80	0.95	1.23	1.01	1.28	1.00	1.07	1.08	1.00	1.08	0.80	0.77	98.0	99.0	0.88	0.72	0.71	69.0	0.64	1.00	0.70	0.75	29.0	0.84	0.55	0.50	0.53	0.37	0.32	0.29
		Grouse	Flushed/Hour	1.05	1.07	1.20	1.13	1.18	1.38	1.42	1.28	0.99	1.10	1.36	1.18	1.35	1.20	0.87	1.04	1.03	1.12	1.08	0.93	1.01	76.0	1.02	0.73	0.68	0.72	0.67	1.06	0.69	0.72	0.55	0.74	0.57	0.47	99.0	0.44	0.40	0.46
		Grouse	Flushed/Trip	4.42	4.62	4.97	5.08	5.19	6.27	6.25	5.43	4.07	4.37	5.55	4.70	5.40	4.87	3.50	4.08	4.09	4.43	4.32	3.79	4.07	3.94	4.06	2.67	2.56	2.69	2.55	3.93	2.69	2.76	1.99	2.62	1.97	1.68	2.30	1.54	1.31	1.56
	# Crouse	Flushed	Reported	5,962	4,553	4,240	4,841	5,618	7,890	8,036	6,749	5,706	5,055	8,653	860,9	868'8	7,071	4,160	5,245	050'5	5,228	4,128	3,212	3,238	2,601	2,855	1,864	1,835	1,483	1,677	3,867	2,475	2,303	1,678	1,842	1,360	1,074	1,167	869	594	650
		Hunting	Hours/Trip	4.16	4.33	4.07	4.16	4.29	4.35	4.34	4.23	4.03	3.91	4.05	3.92	3.96	4.09	4.02	3.88	3.88	3.83	3.86	3.96	3.97	3.92	3.91	3.75	3.66	3.56	3.77	3.72	3.92	3.82	3.60	3.56	3.43	3.58	3.53	3.53	3.28	3.40
			Trips/Hunter	*	17.28	17.79	17.67	15.03	15.54	18.11	18.03	15.74	14.48	19.01	16.06	18.31	16.68	15.06	15.13	14.88	16.40	16.48	16.29	14.20	12.94	13.54	12.45	11.93	9.84	11.15	12.40	12.78	13.22	13.38	11.93	11.16	11.43	9.40	99'6	9.46	8.00
		Respondent	Age	44	44	43	41	43	44	45	45	46	48	49		09	90	51	90	23	53	55	54	54	99	99	55	99	53	54	99	99	99	99	89	69	89	09	09	61	09
ing activ	# H	# Hours	Reported	909'5	4,284	3,474	3,973	4,639	5,479	5,576	5,258	5,642	4,525	6,312	5,042	6,524	5,932	4,788	4,987	4,792	4,519	3,686	3,355	3,160	2,590	2,751	2,616	2,623	1,960	2,482	3,657	3,603	3,186	3,037	2,503	2,375	2,290	1,788	1,605	1,487	1,413
rted hunt	# Hunting	Trips	Reported	1,347	986	854	954	1,082	1,259	1,286	1,244	1,401	1,158	1,559	1,285	1,648	1,451	1,190	1,286	1,235	1,181	926	847	795	099	704	269	716	551	658	983	920	833	843	704	692	640	205	454	454	416
lix I. Repo		# Avid Hunter	Respondents	*	25	48	54	72	81	71	69	89	80	82	80	06	87	79	85	83	72	58	52	99	51	52	56	60	99	69	79	72	63	62	59	62	99	54	47	48	52
Appen			Year	1984-85	1985-86	1986-87	1987-88	1988-89	1989-90	1990-91	1991-92	1992-93	1993-94	1994-95	1995-96	1996-97	1997-98	1998-99	1999-00	2000-01	2001-02	2002-03	2003-04	2004-05	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14	2014-15	2015-16	2016-17	2017-18	2018-19	2019-20	2020-21	2021-22